



Cervical co-infection with high-risk Human Papillomavirus and Herpes simplex

Saadalnour Abusail Mustafa¹, Alfatih Mohamed Ahmed Alnajib², Eldaw Breima Suliman Mohamed^{3,4}, Ahmed Amin Mohammed⁵, Hussain Gadelkarim Ahmed^{6,7}

¹Department of Clinical Laboratory Science, Faculty of Applied Medical Science, Najran University, Najran, Saudi Arabia.

²College of Medicine, University of Ha'il, Saudi Arabia.

³Department of Microbiology, Faculty of Medical Laboratory Science, University of Kordofan, El-Obeid, Sudan.

⁴Shikan College of Medicine, El-Obeid, Sudan.

⁵Department of Obstetrics and Gynecology, Faculty of Medicine, University of Kordofan, El-Obeid, Sudan.

⁶Prof Medical Research consultancy Center, El-Obeid, Sudan.

⁷Department of Histopathology and Cytology, FMLS, University of Khartoum, Sudan.

Abstract

Background: Herpes Simplex Virus (HSV) is a common sexually transmitted virus that infects millions of individuals worldwide. The current study sought to determine the prevalence of Herpes simplex and Human Papillomavirus (HPV) coinfection in Saudi women. **Methodology:** From May 2020 to May 2021, 300 women's cervical smears were collected and sent to a cytopathology laboratory. Because of gynecologic concerns, the women in the study were referred for Pap smears. Cervical materials were then evaluated for the presence of HSV using Polymerase Chain Reaction (PCR) molecular techniques. **Results:** HSV was detected in 2% of patients (66.7% HSV-1 and 33.3% HSV-2). HSV-1 and HPV coinfection was found in 50% of patients, including HPV subtypes 16 and 52. HSV-2 and HPV infection, including HPV subtype 16, was found in 50% of the cases. **Conclusion:** The prevalence of HSV is minimal among Saudi women seeking gynecologic care. Coinfections of HSV and HPV, particularly HPV subtype 16, are prevalent.

Keywords: Herpes simplex virus, Human Papillomavirus, Saudi Arabia, cytology, cervical smear

Correspondence to: Dr. Saadelnour AM, Email: sadosail@yahoo.com

Cite this article: Ammar AB, Eltalib KME, Agab MAAA, Ahmed HG. Medical Research Updates 2023;1(1): 10-22.

Introduction

Herpes simplex virus (HSV) is among the most prevalent sexually transmitted viruses. Infection treatment is not therapeutic, although it can alleviate symptoms and improve quality of life [1]. HSV type 1 or HSV type 2 usually causes genital infection, which can be primary or recurrent. On a regular basis, the virus replicates in epithelial cells and produces latency in sensory neurons before reactivating as a local lesion [2].

HSV is transferred through direct touch, resulting in chronic recurring and painful clinical symptoms. Symptoms include genital infection, cold sores, encephalitis, keratitis, meningitis, and blepharitis.

Cervical cancer and a variety of sexually transmitted diseases can be exacerbated by visible illness [3].

The most common sexually transmitted infection in the world is human papillomavirus (HPV). More than 99% of cervical cancer cases are caused by high-risk HPV subtypes [4]. HSV and HPV coinfection has been reported in gynecologic and colposcopy medical procedures. HSV and high-risk HPV infections have been linked to a history of sexual behavior [5]. HSV type 2 and high-risk HPV coinfection were found to enhance the risk of invasive cervical cancer [6]. In this study, we



looked at Herpes simplex and Human Papillomavirus coinfection in Saudi women.

Materials and Methods

About 300 women were referred to the cytopathology laboratory of the maternity hospital in Al-Madinah, Saudi Arabia, between May 2020 and May 2021. The women in the research were referred for Pap smears due to gynecologic issues. A Pap smear is requested as part of the mandatory investigations in combination with the clinical assessment. Prior to sample collection, each patient was asked to sign a documented ethical consent form.

Cytological samples were obtained by scraping the ectocervix's transition zone. The components were evenly dispersed on a clean glass slide and were promptly fixed in 95% ethyl alcohol while still wet. Following fixation, the smears were stained using the Papanicolaou method (Pap. Method), as follows: the dehydrated smears were soaked in decreasing ethyl alcohol concentrations (90%, 70%, 50%, and finally DW), for two minutes at a time. After five minutes, the smears were stained with Harris' hematoxylin (nuclear stain), rinsed in DW, differentiated in 0.5% aqueous hydrochloric acid for ten seconds (to eliminate remaining stain atoms), and immediately rinsed in DW to end decolorization. The smears were then blued for five seconds in alkaline water before being dehydrated in ethyl alcohol concentrations of 50%, 70%, 90%, and 95% for two minutes each. The smears were then stained for two minutes with Pap. OrangeG6 (cytoplasmic stain), rinsed in 95% ethyl alcohol, and dyed in EA50 (cytoplasmic stain) for three minutes. The smears were then treated with 95% and 100% ethyl alcohol, cleaned in Xylene, and mounted in Distrene Polystyrene Xylene (DPX).

Cervical materials were further tested for the presence of HSV and HPV using molecular methods, such as polymerase chain reaction (PCR), as described by Scouler et al. [7] (See Table 1&2).

DNA extraction

Proteinase K degrades proteins as well as potentially hazardous enzymes such as nucleases. A buffer containing the denaturing agent sodium dodecyl sulfate (SDS) was added to facilitate digestion. Nucleic acids were isolated from tissue lysate using high-speed centrifugation and buffer-

saturated phenol. After that, RNase A was added to the phenol extractions to remove any tainted RNA. Following RNase an incubation, further phenol extractions were performed to remove any remaining enzyme. After adding sodium acetate and isopropanol to precipitate the DNA, high-speed centrifugation was used to pellet the DNA and make it easier to extract the isopropanol. After washing the DNA with 70% ethanol to remove excess salts, it was centrifuged to re-pellet it. DNA is quantified, resuspended in distilled water, and stored at 20 °C. Later, purified DNA was utilized in downstream applications of PCR.

DNA quantification

To assess DNA quantity after DNA extraction, we used a Nano-Drop spectrophotometer to analyze DNA.

Amplification of HPV

Type-specific primers (primer for HPV 31, HPV 33, HPV 35, HPV 39, and HPV 45) were used to detect HPV31, 33, 35, 39, and 45 DNA in cervical lesions using the American Joint Committee on Cancer (AJCC) TNM classification and the International Federation of Gynecology and Obstetrics (FIGO) staging system. The HPV31, 33, 35, 39, and 45 amplification kit from Sacace-Biotechnologies S.r.l. Caserta, Italy was utilized. The total volume of the final reaction was 40 l, which included 10 l of mix-2, 20 l of mix-1 (contained in PCR tubes), and 20 l of extracted DNA (sample). Negative control, positive HPV31,33,35,39, and 45 DNA tubes received 10 l of DNA buffer and 10 l of HPV31,33,35,39, and 45.

Gene Amp PCR system 9700 was used to amp up samples and controls.

Gel-electrophoresis

The PCR products were realized in a 2% Agarose gel with 0.5 g/ml Ethidium bromide. Ten microliters of PCR product and a 100 bp DNA ladder were placed onto the gel. Gel electrophoresis was performed at 120 V and 36 mA for 60 minutes. The photographs were taken with a Gel documentation system (Gel mega, digital camera and computer software).

Table 1. HSV genotype



<i>HSV Genotype</i>	<i>Sequence (5'-3')</i>	<i>Amplification (bp)</i>
<i>HSV-1</i>	CAGACAGCAAAAATCCCCTGAG ACGAGGGAAAACAATAAGG	196
<i>HSV-2</i>	CACCGTCGCCCTATACAGCTT ATCGACGGGATGTGCCAGTTT	210

Table 2 PCR steps. Cycle's time temp PCR steps

<i>Steps</i>	<i>Temperature</i>	<i>Time</i>	<i>Cycles</i>
0	94°C	<i>Pause</i>	
1	94°C	5 min	1
2	94°C	30 sec	25
	48°C	30 sec	
	72°C	25 sec	
3	72°C	1 min	1
4	4°C	<i>Storage</i>	

Statistical Analysis

Data were first organized in an Excel spreadsheet before being moved to SPSS software for analysis to obtain frequencies, percentages, means, cross-tabulations, and the chi-square test. P-values of 0.05 are deemed statistically significant when the 95% confidence interval is used.

Ethical Consent

Before the interview, participants were requested to sign a written ethical consent form. The proposal for the current study was authorized by the Ethical

Committee at the College of Medicine, University Ha'il, Saudi Arabia. 00130/CM-UOH.04/20 HREC.

Results

Three hundred samples from women were examined for the presence of HPV and HSV. The women's ages ranged from 20 to 70. As shown in Fig 1, the majority of women were between the ages of 31 and 40, with 41-50, 51-60, 20-30, and >60 years constituting 27.7%, 26.7%, 24.3%, 10.7%, and 10.7%, respectively.




Figure 1. Description of the study subjects by age.

HSV was discovered positive in 6/300 (2%) of the patients in this series (4/6(66.7%) HSV-1 and 2/6(33.3%) HSV-2). HSV-1 and HPV co-infection was found in 2/4 (50%) of the patients, which

comprised HPV subtypes 16 and 52. HSV-2 and HPV coinfection was found in 12 (50%) of the cases, including HPV subtype 16, as shown in Table 1 and Figure 2.

Table 1. Correlation between HSV and HPV

Variable	Herpes Simples				Total
	HSV-1 Positive	HSV-1 Negative	HSV-1 Total	HSV-2 Positive	
<i>HPV</i>					
Positive	2	12	14	1	14
Negative	2	284	286	1	286
Total	4	296	300	2	300
<i>HPV-Subtypes</i>					
HPV-16	1	5	6	1	6
HPV-18	0	1	1	0	1
HPV-52	1	2	3	0	3
HPV-58	0	2	2	0	2
HPV-33	0	2	2	0	2
Total	2	12	14	1	14




Figure 2. HSV and HPV co-infection

Atypical cytological alterations were found in 4/6 (67%) of HSV-positive individuals, including 2/4 (50%) of HSV-1 cases and 2/2 (100%) of HSV-2 cases. In two cases, inflammatory cell infiltrates

were observed: one (25%) HSV-1 and one (50%) HSV-2. As shown in Table 2 Fig 3, all Candida Albicans cases tested negative for HSV.

Table 2. Distribution of HSV by cervical cytopathology

Variable	Herpes simplex				Total
	HSV-1 Positive	HSV-1 Negative	HSV-2 Positive	HSV-2 Negative	
Cytological atypia					
Normal	2	265	1	266	267
Mild atypia	0	10	1	10	11
Moderate	0	11	0	10	10
Severe	2	10	0	12	12
Total	4	296	2	298	300
Inflammatory change					
Yes	1	70	1	70	71
No	3	226	1	228	229
Total	4	296	2	298	300
Candida Albicans					
Yes	0	7	0	7	7
No	4	289	2	291	293
Total	4	296	2	298	300




Figure 3. HSV by cervical cytopathology

As shown in Table 3, approximately 2/4 (50%) of the HSV-1 positive cases were seen in the age group >60 years, one (25%) in the age group 51-60 years, and one (25%) in the age group 41-50 years,

whereas one (50%) case HSV-2 was seen in the age group 20-30 years and the remaining one in the age group 41-50 years.

Table 3. HSV infection status by age

Variable	Herpes simplex				Total
	HSV-1		HSV-2		
	Positive	Negative	Positive	Negative	
20-30 years	0	32	1	31	32
31-40	0	83	0	83	83
41-50	1	79	1	79	80
51-60	1	72	0	73	73
>60	2	30	0	32	32
Total	4	296	2	298	300



Figure 4. PCR amplification of HSV-1 and HSV-2 in cervical samples.

The products were separated on a 2% agarose gel and stained with an ethidium bromide 1000bp ladder (arrows show 100 and 200 bands), with HSV-1 positive in 196 bp. HSV-2 is not present.

Discussion

In this Saudi patient series, the overall HSV prevalence was 2% (with roughly 67% HSV type 1 and approximately 33% HSV type 2). Saudi Arabian epidemiological data on HSV are limited. A Saudi Arabian study assessed IgG and IgM antibodies in women and neonates in a group of patients with recurrent abortion. The presence of IgM in 1.3% of women indicated a recent HSV type 1 infection. The presence of IgG, observed in around 81% of the women, indicated past HSV exposure.

As a result, HSV infection is associated with infertility [8]. Another Saudi Arabian study included 4985 women who were screened for HSV.

The total prevalence of HSV type 1 was 88.8%, while HSV type 2 was 1.26%, according to the study. The incidence was substantially higher among those who were married, divorced, or widowed. Other sexually transmitted diseases, such as *Treponema pallidum* and the Human Immunodeficiency Virus (HIV), have been found to be more common [9]. However, 491.5 persons were predicted to be infected with HSV type 2, accounting for 13.2% of the world's population aged 15-49. Furthermore, 3752 million people were found to be infected with HSV type 1, for a global prevalence of 66.6%. Infection patterns



varied according to geographical area, sex, and age [10].

In the current study, almost half of the cases of HSV infection had HPV coinfection. Some investigations have assumed HSV and HPV virus co-infection, as well as other carcinogenic viruses [11,12]. Furthermore, some researchers believe that HSV infection may encourage subsequent HPV infection [13]. Coinfection occurred with a variety of HPV subtypes, with a higher prevalence of HPV subtype 16.

The current study found that HSV type 1 was linked to severe cytological atypia, whereas HSV type 2 was linked to mild cytological atypia. However, HSV replication in human epithelial cells causes structural alterations in the cell cytoplasmic organelles and nucleus, which eventually result in visible morphological abnormalities [14]. Infection with HSV type 1 disrupts the end of transcription of RNA polymerase II in host genes [15].

In the current investigation, inflammatory cells were found in two cases, one (25%) HSV-1 and one (50%) HSV-2.

HSV can activate both innate and adaptive immune responses in host cells, triggering a cascade of

Authors contribution

SAM: Conceptual, consultation, funding, and approval of the final version

AMAA: Conceptual, data analysis, funding, and approval of the final version

EBSM: conceptual, manuscript drafting, and approval of the final version

AAM: conceptual, administration, funding, and approval of the final version

HGA: conceptual, administration, funding, and approval of the final version

events that results in the release of inflammatory mediators that recruit a large number of inflammatory cells [16].

In the current study, HSV infections were more common in older adults (> 50 years old); the incidence of HSV by age varies according to a variety of factors, including geographical region, community sexual behavior, and population subgroups. Females are more likely than males to be infected with HSV type 2 [17].

The current analysis revealed some hidden HSV data in Saudi Arabia, but it has significant limitations, including a lack of previous immune globulins, which predicts previous exposure.

In conclusion: The frequency of HSV is low among Saudi women seeking gynecologic care. Coinfections of HSV and HPV, particularly HPV subtype 16, are prevalent.

Acknowledgment

The authors would like to thank Dr. Hashim M. Missawi at the Department of clinical pathology, maternity, and children's hospital, for his help in sample collection and consultation.

Funding

Self-funded.

Data availability

The participants of this study did not give written consent for their data to be shared publicly, so due to the sensitive nature of the research supporting data is not available.

Disclosure of interest

No interest to declare

References

- 1- Cole S. Herpes Simplex Virus: Epidemiology, Diagnosis, and Treatment. *Nurs Clin North Am.* 2020 Sep;55(3):337-345. doi: 10.1016/j.cnur.2020.05.004.
- 2- Mathew Jr J, Sapra A. Herpes Simplex Type 2. 2021 Feb 23. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. PMID: 32119314.
- 3- Caputo A, Marconi P. Vaccine Development for Herpes Simplex Viruses: A Commentary of Special Issue Editors. *Vaccines (Basel).* 2021 Feb 16;9(2):158. doi: 10.3390/vaccines9020158. PMID: 33669413; PMCID: PMC7920455.
- 4- Yuan Y, Cai X, Shen F, Ma F. HPV post-infection microenvironment and cervical cancer. *Cancer Lett.* 2021 Jan 28; 497:243-254. doi: 10.1016/j.canlet.2020.10.034.
- 5- Bahena-Román M, Sánchez-Alemán MA, Contreras-Ochoa CO, Lagunas-Martínez A, Olamendi-Portugal M, López-Estrada G, Delgado-Romero K, Guzmán-Olea E, Madrid-Marina V, Torres-Poveda K. Prevalence of active infection by herpes simplex virus type 2 in patients with high-risk human papillomavirus infection: A cross-sectional study. *J Med Virol.* 2020 Aug;92(8):1246-1252. doi: 10.1002/jmv.25668.
- 6- Smith JS, Herrero R, Bosetti C, Muñoz N, Bosch FX, Eluf-Neto J, Castellsagué X, Meijer CJ, Van den Brule AJ, Franceschi S, Ashley R; International Agency for Research on Cancer (IARC) Multicentric Cervical Cancer Study Group. Herpes simplex virus-2 as a human papillomavirus cofactor in the etiology of invasive cervical cancer. *J Natl Cancer Inst.* 2002 Nov 6;94(21):1604-13. doi: 10.1093/jnci/94.21.1604.
- 7- Scoular A, Gillespie G, Carman WF. Polymerase chain reaction for diagnosis of genital herpes in a genitourinary medicine clinic. *Sex Transm Infect.* 2002 Feb;78(1):21-5. doi: 10.1136/sti.78.1.21.
- 8- Althaqafi RMM, Elrewiny M, Abdel-Moneim AS. Maternal and neonatal infections of herpes simplex virus-1 and cytomegalovirus in Saudi Arabia. *J Infect Public Health.* 2020 Feb;13(2):313-314. doi: 10.1016/j.jiph.2019.11.020.
- 9- Memish ZA, Almasri M, Chentoufi AA, Al-Tawfiq JA, Al-Shangiti AM, Al-Kabbani KM, Otaibi B, Assirri A, Yezli S. Seroprevalence of Herpes Simplex Virus Type 1 and Type 2 and Coinfection with HIV and Syphilis: The First National Seroprevalence Survey in Saudi Arabia. *Sex Transm Dis.* 2015 Sep;42(9):526-32. doi: 10.1097/OLQ.0000000000000336. PMID: 26267880.
- 10- James C, Harfouche M, Welton NJ, Turner KM, Abu-Raddad LJ, Gottlieb SL, Looker KJ. Herpes simplex virus: global infection prevalence and incidence estimates, 2016. *Bull World Health Organ.* 2020 May 1;98(5):315-329. doi: 10.2471/BLT.19.237149.
- 11- Guidry JT, Scott RS. The interaction between human papillomavirus and other viruses. *Virus Res.* 2017; 231:139-147. doi: 10.1016/j.virusres.2016.11.002.
- 12- Akbari M, Elmi R. Herpes Simplex Virus and Human Papillomavirus Coinfections in Hyperimmunoglobulin E Syndrome Presenting as a Conjunctival Mass Lesion. *Case Rep Med.* 2017; 2017:1650841. doi: 10.1155/2017/1650841. Epub 2017 Oct 12. PMID: 29158737; PMCID: PMC5660752.
- 13- Mäki J, Paavilainen H, Kero K, Hukkanen V, Syrjänen S. Herpes simplex and human papilloma virus coinfections in oral mucosa of men-A 6-year follow-up study. *J Med Virol.* 2018 Mar;90(3):564-570. doi: 10.1002/jmv.24965.
- 14- Ahmad I, Wilson DW. HSV-1 Cytoplasmic Envelopment and Egress. *Int J Mol Sci.* 2020 Aug 19;21(17):5969. doi: 10.3390/ijms21175969.
- 15- Wang X, Hennig T, Whisnant AW, Erhard F, Prusty BK, Friedel CC, Forouzmand E, Hu W, Erber L, Chen Y, Sandri-Goldin RM, Dölken L, Shi Y. Herpes simplex virus blocks host transcription termination via the bimodal activities of ICP27. *Nat Commun.* 2020 Jan 15;11(1):293. doi: 10.1038/s41467-019-14109-x.
- 16- Wang L, Wang R, Xu C, Zhou H. Pathogenesis of Herpes Stromal Keratitis: Immune Inflammatory Response Mediated by Inflammatory Regulators. *Front Immunol.* 2020 May 13; 11:766. doi: 10.3389/fimmu.2020.00766.
- 17- Smith JS, Robinson NJ. Age-specific prevalence of infection with herpes simplex virus types 2 and 1: a global review. *J Infect Dis.* 2002 Oct 15;186 Suppl 1: S3-28. doi: 10.1086/343739.